Few-Body Systems manuscript No.
(will be inserted by the editor)

Sergio Scopetta · Alessio Del Dotto · Leonid Kaptari · Emanuele Pace · Matteo Rinaldi · Giovanni Salmè

A Light-Front approach to the ^{3}He spectral function

arXiv:1411.7559v1 [nucl-th] 27 Nov 2014

Received: date / Accepted: date

Abstract The analysis of semi-inclusive deep inelastic electron scattering off polarized ^{3}He at finite momentum transfers, aimed at the extraction of the quark transverse-momentum distributions in the neutron, requires the use of a distorted spin-dependent spectral function for ^{3}He, which takes care of the final state interaction effects. This quantity is introduced in the non-relativistic case, and its generalization in a Poincaré covariant framework, in plane wave impulse approximation for the moment being, is outlined. Studying the light-front spin-dependent spectral function for a J=1/2 system, such as the nucleon, it is found that, within the light-front dynamics with a fixed number of constituents and in the valence approximation, only three of the six leading twist T-even transverse-momentum distributions are independent.

1 Introduction

Information on the three-dimensional proton structure can be obtained from the quark transverse momentum distributions (TMDs) [1], which can be accessed through semi-inclusive deep inelastic electron scattering (SIDIS). In particular the single spin asymmetries (SSAs) allow one to extract the Sivers and the Collins contributions, expressed in terms of different TMDs and fragmentation functions (ff) [1]. A large Sivers asymmetry was measured in $\mathbf{p}(e,e'\pi)x$ [2] and a small one in $\mathbf{D}(e,e'\pi)x$ [3], showing a strong flavor dependence of TMDs. To clarify the flavour dependence of the quark transverse momentum distributions, high precision experiments, involving both protons and neutrons, are needed [4].

In Ref. [5] an experiment to extract information on the neutron TMDs from experimental measurements of the SSAs on ^{3}He, at JLab at 12 GeV, was proposed. To obtain a reliable information one has

S. Scopetta
Dipartimento di Fisica e Geologia, Università di Perugia and INFN, Sezione di Perugia, Italy
Tel.: +39-075-5852721
Fax: +39-075-44666
E-mail: sergio.scopetta@pg.infn.it

A. Del Dotto
Dipartimento di Fisica, Università di Roma Tre and INFN, Roma 3, Italy

L. Kaptari
Bogoliubov Lab. Theor. Phys., 141980, JINR, Dubna, Russia

E. Pace
Dipartimento di Fisica, Università di Roma "Tor Vergata" and INFN, Roma 2, Italy

M. Rinaldi
Dipartimento di Fisica e Geologia, Università di Perugia and INFN, Sezione di Perugia, Italy

G. Salmè
INFN Sezione di Roma, Italy

to take carefully into account the structure of ^{3}He, the interaction in the final state (FSI) between the observed pion and the remnant debris, and the relativistic effects. The present paper reports on our efforts about these items.

2 SIDIS off ^{3}He

A polarized ^{3}He is an ideal target to study the polarized neutron since, at a 90% level, a polarized ^{3}He is equivalent to a polarized neutron. Dynamical nuclear effects in inclusive deep inelastic electron scattering (DIS) $^3\mathbf{He}(e,e')X$ (DIS) were evaluated [6] with a realistic $^3\mathbf{He}$ spin-dependent spectral function, $P^{\tau}_{\sigma,\sigma\prime}(\mathbf{p},E,S_{He})$, with $\mathbf{p}$ the initial nucleon momentum in the laboratory frame and E the missing energy [7]. It was found that the formula

$$A_n \simeq \frac{1}{p_n f_n}\left(A_3^{exp} - 2p_p f_p\ A_p^{exp}\right) \tag{1}$$

can be safely adopted to extract the neutron information from ^{3}He and proton data. This formula is actually widely used by experimental collaborations. The nuclear effects are hidden in the proton and neutron "effective polarizations", $p_{p(n)}$. With the AV18 nucleon-nucleon interaction [8] the values $p_p = -0.023$, $p_n = 0.878$ were obtained [9]. The quantities $f_{p(n)}$ in Eq. (1) are the dilution factors. To investigate if an analogous formula can be used to extract the SSAs, in [9] the processes $^3\mathbf{He}(e,e'\pi^{\pm})X$, with ^{3}He transversely polarized, were evaluated in the Bjorken limit and in PWIA, i.e. the final state interaction (FSI) was considered only within the two-nucleon spectator pair which recoils. In such a framework, SSAs for ^{3}He involve convolutions of $P^{\tau}_{\sigma,\sigma\prime}(\mathbf{p},E,S_{He})$, with TMDs and ffs. Ingredients of the calculations were: i) a realistic $P^{\tau}_{\sigma,\sigma\prime}(\mathbf{p},E,S_{He})$ for ^{3}He, obtained using the AV18 interaction; ii) parametrizations of data for TMDs and ff, whenever available; iii) models for the unknown TMDs and ff. It was found that, in the Bjorken limit, the extraction procedure through the formula successful in DIS works nicely for the Sivers and Collins SSAs [9]. The generalization of Eq. (1) to extract Sivers and Collins asymmetries from ^{3}He and proton asymmetries was recently used by experimental collaborations [10].

In SIDIS experiments off ^{3}He, the relative energy between the spectator $(A-1)$ system and the system composed by the detected pion and the remnant debris (see Fig. 1) is a few GeV and FSI can be treated through a generalized eikonal approximation (GEA) [11]. The GEA was already succesfully applied to unpolarized SIDIS in Ref. [12]. The FSI effects to be considered are due to the propagation of the debris, formed after the γ^* absorption by a target quark, and the subsequent hadronization, both of them influenced by the presence of a fully interacting $(A-1)$ spectator system (see Fig. 1).



Fig. 1 Interaction between the $(A-1)$ spectator system and the debris produced by the absorption of a virtual photon by a nucleon in the nucleus.

In this approach, the key quantity to introduce the FSI effects is the *distorted* spin-dependent spectral function, whose relevant part in the evaluation of SSAs is:

$$\mathcal{P}_{||}^{PWIA(FSI)} = \mathcal{O}_{\frac{1}{2}\frac{1}{2}}^{IA(FSI)} - \mathcal{O}_{-\frac{1}{2}-\frac{1}{2}}^{IA(FSI)} \tag{2}$$

with:

$$\mathcal{O}_{\lambda\lambda'}^{IA}(\mathbf{p},E) = \sum\!\!\!\!\!\!\int d\epsilon^*_{A-1}\ \rho\left(\epsilon^*_{A-1}\right)\ \langle S_A,\mathbf{P}_A|\Phi_{\epsilon^*_{A-1}},\lambda',\mathbf{p}\rangle\langle\Phi_{\epsilon^*_{A-1}},\lambda,\mathbf{p}|S_A,\mathbf{P}_A\rangle\ \delta\left(E-B_A-\epsilon^*_{A-1}\right)\ , \tag{3}$$

and

$$\begin{aligned}\mathcal{O}_{\lambda\lambda'}^{FSI}(\mathbf{p},E) = \sum\!\!\!\!\!\!\int d\epsilon^*_{A-1}\ \rho\left(\epsilon^*_{A-1}\right)\ &\langle S_A,\mathbf{P}_A|(\hat{S}_{Gl})\{\Phi_{\epsilon^*_{A-1}},\lambda',\mathbf{p}\}\rangle\langle(\hat{S}_{Gl})\{\Phi_{\epsilon^*_{A-1}},\lambda,\mathbf{p}\}|S_A,\mathbf{P}_A\rangle\\ &\times\ \delta\left(E-B_A-\epsilon^*_{A-1}\right)\ ,\end{aligned} \tag{4}$$

where $\rho\left(\epsilon^*_{A-1}\right)$ is the density of the $(A-1)$-system states with intrinsic energy ϵ^*_{A-1}, and $|S_A,\mathbf{P}_A\rangle$ is the ground state of the A-nucleons nucleus with polarization S_A. $\hat{S}_{Gl}$ is the Glauber operator:

$$\hat{S}_{Gl}(\mathbf{r}_1,\mathbf{r}_2,\mathbf{r}_3) = \prod_{i=2,3}\left[1-\theta(z_i-z_1)\Gamma(\mathbf{b}_1-\mathbf{b}_i,z_1-z_i)\right] \tag{5}$$

and $\Gamma(\mathbf{b}_{1i},z_{1i})$ the generalized profile function:

$$\Gamma(\mathbf{b}_{1i},z_{1i})\ =\ \frac{(1-i\,\alpha)\ \sigma_{eff}(z_{1i})}{4\,\pi\,b_0^2}\ \exp\left[-\frac{\mathbf{b}_{1i}^2}{2\,b_0^2}\right]\ , \tag{6}$$

where $\mathbf{r}_{1i}=\{\mathbf{b}_{1i},\mathbf{z}_{1i}\}$ with $\mathbf{z}_{1i}=\mathbf{z}_1-\mathbf{z}_i$ and $\mathbf{b}_{1i}=\mathbf{b}_1-\mathbf{b}_i$. The models for the profile function, $\Gamma(\mathbf{b}_{1i},z_{1i})$, and for the effective cross section, $\sigma_{eff}(z_{1i})$, as well as the values of the parameters α and b_0 are the ones exploited in Ref. [12] to nicely describe the JLab data corresponding to the unpolarized spectator SIDIS off the deuteron.

It occurs that $\mathcal{P}_{||}^{PWIA}$ and $\mathcal{P}_{||}^{FSI}$ can be very different, but the relevant observables for the SSAs involve integrals, dominated by the low momentum region, where the FSI effects on $\mathcal{P}_{||}$ are minimized and the spectral function is large [11]. As a consequence the effective nucleon polarizations change from $p_p=-0.023$, $p_n=0.878$ to $p_p^{FSI}=-0.026$, $p_n^{FSI}=0.76$, where

$$p_{p(n)}^{FSI} = \int d\epsilon_S \int d\mathbf{p}\ Tr[\mathbf{S}_{He}*\sigma\ P_{FSI}^{p(n)}(\mathbf{p},E,S_{He})]\ , \tag{7}$$

with $P_{FSI}^{p(n)}(\mathbf{p},E,S_{He})$ the distorted spin-dependent spectral function, defined in terms of the overlaps of Eq. (4) [11]. Then $p_{p(n)}$ with and without the FSI differ by 10-15% . Actually, one has also to consider the effect of the FSI on dilution factors. We have found, in a wide range of kinematics typical for the experiments at JLAB [5; 10], that the product of polarizations and dilution factors changes very little [13]. Indeed the effects of FSI in the dilution factors and in the effective polarizations are found to compensate each other to a large extent and the usual extraction appears to be safe :

$$A_n \simeq \frac{1}{p_n^{FSI}f_n^{FSI}}\left(A_3^{exp}-2p_p^{FSI}f_p^{FSI}A_p^{exp}\right)\ \simeq\frac{1}{p_nf_n}\left(A_3^{exp}-2p_pf_p\ A_p^{exp}\right) \tag{8}$$

In [9] the calculation was performed within a non relativistic approach for the spectral function, but with the correct relativistic kinematics in the Bjorken limit. For an accurate description of SIDIS processes, the role played by relativity has to be fully investigated: it will become even more important with the upgrade of JLab @ 12 GeV. To study relativistic effects in the actual experimental kinematics, we adopted [14] the Relativistic Hamiltonian Dynamics (RHD) introduced by Dirac [15]. Indeed the RHD of an interacting system with a fixed number of on-mass-shell constituents, with the Bakamijan-Thomas construction of the Poincaré generators [16], is fully Poincaré covariant. The Light-Front (LF) form of RHD has a subgroup structure of the LF boosts, allows a separation of the intrinsic motion from the global one, which is very important for the description of DIS, SIDIS and deeply virtual Compton scattering (DVCS) processes, and allows a meaningful Fock expansion. The key quantity to consider in SIDIS processes is the LF relativistic spectral function, $\mathcal{P}^{\tau}_{\sigma'\sigma}(\tilde{\kappa},\epsilon_S,S_{He})$, with $\tilde{\kappa}$ an intrinsic

Table 1 Proton and neutron effective polarizations within the non relativistic appproach (NR) and preliminary results within the light-front relativistic dynamics approach (LF). First line : longitudinal effective polarizations; second line : transverse effective polarizations.

	$proton\,NR$	$proton\,LF$	$neutron\,NR$	$neutron\,LF$
$\int dEd\mathbf{p}\ Tr(\mathcal{P}\sigma_z)_{\mathbf{S}_A=\widehat{z}}$	-0.02263	-0.02231	0.87805	0.87248
$\int dEd\mathbf{p}\ Tr(\mathcal{P}\sigma_y)_{\mathbf{S}_A=\widehat{y}}$	-0.02263	-0.02268	0.87805	0.87494

nucleon momentum and ϵ_S the energy of the two-nucleon spectator system. The LF spectral function will be very useful also for other studies (e.g., for nuclear generalized parton distributions (GPDs), where final states have to be properly boosted, studied so far only within a non-relativistic spectral function [17; 18]). The LF nuclear spectral function, $\mathcal{P}^{\tau}_{\sigma'\sigma}(\tilde{\kappa},\epsilon_S,S_{He})$, is defined in terms of LF overlaps [19] between the ground state of a polarized ^{3}He and the cartesian product of an interacting state of two nucleons with energy ϵ_S and a plane wave for the third nucleon. Within a reliable approximation [19] it can be given in terms of the unitary Melosh Rotations, $D^{\frac{1}{2}}[\mathcal{R}_M(\tilde{\kappa})]$, and the usual instant-form spectral function $P^{\tau}_{\sigma'_1\sigma_1}$:

$$\mathcal{P}^{\tau}_{\sigma'\sigma}(\tilde{\kappa},\epsilon_S,S_{He}) \propto \sum_{\sigma_1\sigma'_1} D^{\frac{1}{2}}[\mathcal{R}^{\dagger}_M(\tilde{\kappa})]_{\sigma'\sigma'_1}\ P^{\tau}_{\sigma'_1\sigma_1}(\mathbf{p},\epsilon_S,S_{He})\ D^{\frac{1}{2}}[\mathcal{R}_M(\tilde{\kappa})]_{\sigma_1\sigma} \tag{9}$$

Notice that the instant-form spectral funtion $P^{\tau}_{\sigma'_1\sigma_1}(\mathbf{p},\epsilon_S,S_{He})$ is given in terms of three independent functions, B_0, B_1, B_2 [7], once parity and t-reversal are imposed:

$$P^{\tau}_{\sigma'_1\sigma_1}(\mathbf{p},\epsilon_S,S_{He}) = \left[B^{\tau}_{0,S_{He}}(|\mathbf{p}|,E) + \boldsymbol{\sigma}\cdot\mathbf{f}^{\tau}_{S_{He}}(\mathbf{p},E)\right]_{\sigma'_1\sigma_1} \tag{10}$$

with

$$\mathbf{f}^{\tau}_{S_{He}}(\mathbf{p},E) = \mathbf{S}_{He}B^{\tau}_{1,S_{He}}(|\mathbf{p}|,E) + \hat{\mathbf{p}}\ (\hat{\mathbf{p}}\cdot\mathbf{S}_{He})B^{\tau}_{2,S_{He}}(|\mathbf{p}|,E) \tag{11}$$

Adding FSI, more terms should be included.

We are now evaluating the SSAs using the LF hadronic tensor, at finite values of Q^2. The preliminary results are quite encouraging, since, as shown in Table 1, LF longitudinal and transverse polarizations change little from the ones obtained within the NR spectral function and weakly differ from each other. Then we find that the extraction procedure works well within the LF approach as it does in the non relativistic case.

Concerning the FSI, we plan to include in our LF approach the FSI between the jet produced from the hadronizing quark and the two-nucleon system through the Glauber approach of Ref. [11].

3 The $J = 1/2$ LF spectral function and the nucleon LF TMDs

The TMDs for a nucleon with total momentum P and spin S are introduced through the q-q correlator

$$\begin{aligned}\Phi(k,P,S)_{\alpha\beta} = \int d^4z\ e^{ik\cdot z}\langle PS|\ \bar{\psi}_{q\beta}(0)\ \psi_{q\alpha}(z)|PS\rangle = \frac{1}{2}\{\ A_1\ \not{P}\ + A_2\ S_L\ \gamma_5\ \not{P}\ + A_3\ \not{P}\gamma_5\ S_\perp \\ +\ \frac{1}{M}\ \widetilde{A}_1\ \mathbf{k}_\perp\cdot\mathbf{S}_\perp\ \gamma_5\not{P}\ +\ \widetilde{A}_2\ \frac{S_L}{M}\ \not{P}\gamma_5\ \not{k}_\perp\ \frac{1}{M^2}\ \widetilde{A}_3\ \mathbf{k}_\perp\cdot\mathbf{S}_\perp\ \not{P}\gamma_5\ \not{k}_\perp\}_{\alpha\beta},\end{aligned} \tag{12}$$

where k is the parton momentum in the laboratory frame, so that the six twist-2 T-even functions, $A_i,\ \widetilde{A}_i\ (i=1,3)$, can be obtained by proper traces of $\Phi(k,P,S)$.

Indeed, particular combinations of the functions $A_i, \widetilde{A}_i\ (i=1,3)$ can be obtained by the following traces of $\Phi(k,P,S)$:

$$\begin{aligned}\frac{1}{2P^+}\,\mathrm{Tr}(\gamma^+\Phi) &= A_1\,,\\ \frac{1}{2P^+}\,\mathrm{Tr}(\gamma^+\gamma_5\Phi) &= S_L\,A_2\ +\ \frac{1}{M}\,\mathbf{k}_\perp\cdot\mathbf{S}_\perp\,\widetilde{A}_1\,,\\ \frac{1}{2P^+}\,\mathrm{Tr}(\mathrm{i}\sigma^{\mathrm{j}+}\gamma_5\Phi) &= S^j_\perp\,A_3\ +\ \frac{S_L}{M}\,k^j_\perp\,\widetilde{A}_2\ +\frac{1}{M^2}\ \mathbf{k}_\perp\cdot\mathbf{S}_\perp\ k^j_\perp\ \widetilde{A}_3\quad (j=x,y).\end{aligned} \tag{13}$$

The six T-even twist-2 TMDs for the quarks inside a nucleon can be obtained by integration of the functions $A_i,\ \widetilde{A}_i$ on k^+ and k^- as follows

$$f(x,|\mathbf{k}_\perp|^2) = \frac{1}{2}\int \frac{dk^+dk^-}{(2\pi)^4}\ \delta[k^+ - xP^+]\ 2P^+ A_1\,,$$
$$\Delta f(x,|\mathbf{k}_\perp|^2) = \frac{1}{2}\int \frac{dk^+dk^-}{(2\pi)^4}\ \delta[k^+ - xP^+]\ 2P^+ A_2\,,$$
$$g_{1T}(x,|\mathbf{k}_\perp|^2) = \frac{1}{2}\int \frac{dk^+dk^-}{(2\pi)^4}\ \delta[k^+ - xP^+]\ 2P^+ \widetilde{A}_1\,,$$
$$\Delta'_T f(x,|\mathbf{k}_\perp|^2) = \frac{1}{2}\int \frac{dk^+dk^-}{(2\pi)^4}\ \delta[k^+ - xP^+]\ 2P^+ \left(A_3 + \frac{|\mathbf{k}_\perp|^2}{2M^2}\widetilde{A}_3\right)\,,$$
$$h_{1L}^\perp(x,|\mathbf{k}_\perp|^2) = \frac{1}{2}\int \frac{dk^+dk^-}{(2\pi)^4}\ \delta[k^+ - xP^+]\ 2P^+ \widetilde{A}_2\,,$$

$$h_{1T}^\perp(x,|\mathbf{k}_\perp|^2) = \frac{1}{2}\int \frac{dk^+dk^-}{(2\pi)^4}\ \delta[k^+ - xP^+]\ 2P^+ \widetilde{A}_3\ . \tag{14}$$

Let us consider the contribution to the correlation function from on-mass-shell fermions

$$\Phi_p(k,P,S) = \frac{(\not{k}_{on} + m)}{2m}\ \Phi(k,P,S)\ \frac{(\not{k}_{on} + m)}{2m} = \tag{15}$$

$$= \sum_{\sigma\sigma'}\ u_{LF}(\tilde{k},\sigma')\ \bar{u}_{LF}(\tilde{k},\sigma')\ \Phi(k,P,S)\ u_{LF}(\tilde{k},\sigma)\bar{u}_{LF}(\tilde{k},\sigma)\ ,$$

and let us identify $\bar{u}_{LF}(\tilde{k},\sigma')\ \Phi(k,P,S)\ u_{LF}(\tilde{k},\sigma)$, up to a kinematical factor K, with the LF nucleon spectral function, $\mathcal{P}_{\sigma'\sigma}(\tilde{\kappa},\epsilon_S,S)$ [19]:

$$\bar{u}_{LF}(\tilde{k},\sigma')\ \Phi(k,P,S)\ u_{LF}(\tilde{k},\sigma) = K\ \mathcal{P}_{\sigma'\sigma}(\tilde{\kappa},\epsilon_S,S)\ . \tag{16}$$

In a reference frame where $\mathbf{P}_\perp = 0$, the following relation holds between the off-mass-shell minus component k^- of the momentum of the struck quark and the spectator diquark energy ϵ_S :

$$k^- = \frac{M^2}{P^+} - \frac{(\epsilon_S + m)\ 4m + |\boldsymbol{k}_\perp|^2}{P^+ - k^+}\ . \tag{17}$$

Let us approximate the full correlation function $\Phi(k,P,S)$ by its particle contribution $\Phi_p(k,P,S)$. Then, through relations analogous to the ones of Eq. (13), which allow one to obtain the functions $A_i,\ \widetilde{A}_i$ from the traces of $\Phi(k,P,S)$, the valence approximations $A_i^V,\ \widetilde{A}_i^V\ (i=1,3)$ for the functions $A_i,\ \widetilde{A}_i$ can be obtained by the traces $[\gamma^+\ \Phi_p(k,P,S)]$, $[\gamma^+\ \gamma_5\ \Phi_p(k,P,S)]$, and $[\not{k}_\perp\gamma^+\gamma_5\ \Phi_p(k,P,S)]$. However these same traces can be also expressed through the LF spectral function, since from Eqs. (15,16) one has

$$Tr\left[\gamma^+\ \Phi_p(k,P,S)\right] = \frac{k^+}{m}\ K\ Tr\left[\mathcal{P}(\tilde{\kappa},\epsilon_S,S)\right] \tag{18}$$
$$Tr\left[\gamma^+\ \gamma_5\ \Phi_p(k,P,S)\right] = \frac{k^+}{m}\ K\ Tr\left[\sigma_z\ \mathcal{P}(\tilde{\kappa},\epsilon_S,S)\right] \tag{19}$$
$$Tr\left[\not{k}_\perp\gamma^+\gamma_5\ \Phi_p(k,P,S)\right] = \frac{k^+}{m}KTr\left[\boldsymbol{k}_\perp\cdot\boldsymbol{\sigma}\ \mathcal{P}(\tilde{\kappa},\epsilon_S,S)\right] \tag{20}$$

In turn, as in the ^{3}He case, the traces $\mathrm{Tr}(\mathcal{P}I)$, $\mathrm{Tr}(\mathcal{P}\sigma_z)$, $\mathrm{Tr}(\mathcal{P}\sigma_i)$ $(i=x,y)$ can be expressed in terms of three scalar functions, B_0, B_1, B_2 and known kinematical factors. Then, within the LF approach with a fixed number of particles and in the valence approximations, the six leading twist T-even functions $A_i^V,\ \widetilde{A}_i^V\ (i=1,3)$ can be expressed in terms of the three independent scalar functions B_0, B_1, B_2. Therefore only three of the six T-even TMDs are independent [19]. We stress however that this result, not valid in general in QCD (see, e.g., Ref. [20]), is a prediction of our peculiar framework, i.e., hamiltonian light-front dynamics with a fixed number of constituents, finalized to yield a proper Poincaré covariant description of the nucleon in the valence approximation. Its experimental check would be therefore a test of the reliability of our scheme to describe SIDIS processes in the valence region.

4 Conclusions

A realistic study of the DIS processes off ^{3}**He** and in particular of the SSAs in the reaction $^3\mathbf{He}(e, e'\pi^{\pm})X$ beyond the PWIA and the non relativistic approach is under way. The FSI effects have been evaluated through the GEA and the introduction of a distorted spin-dependent spectral function. The relativistic effects are studied through the analysis of a LF spectral function (up to now only in PWIA). Preliminary results are encouraging, in view of a sound extraction of the neutron information from experimental data. Nuclear effects in the extraction of the neutron information are found to be under control, even when the interaction in the final state is considered, and the relativistic effects appear to be small. An analysis at finite Q^2 with the LF spectral function is in progress. The next step is to complete this program. Then we will apply the LF spectral function to other processes (e.g., DVCS) to exploit other possibilities to use ^{3}He as an effective neutron target and as a laboratory for light-front studies.

The introduction of a LF spin-dependent spectral function for a nucleon allowed us to find relations among the six leading twist T-even TMDs, which are exact within LF dynamics with a fixed number of degrees of freedom, in the valence approximation. It can be shown that, in this case, only three of the six T-even TMDs are independent. The above relations, although not true in QCD, could be experimentally checked to test our LF description of SIDIS in the valence region.

References

1. Barone, V., Drago, A., and Ratcliffe, P. G., "Transverse polarisation of quarks in hadrons," Phys. Rept. **359**, 1 (2002).
2. Airapetian, A. *et al.* [HERMES Collaboration], "Single-spin asymmetries in semi-inclusive deep-inelastic scattering on a transversely polarized hydrogen target," Phys. Rev. Lett. **94**, 012002 (2005) .
3. Alexakhin, V. Y *et al.* [COMPASS Collaboration], "First measurement of the transverse spin asymmetries of the deuteron in semi-inclusive deep inelastic scattering," Phys. Rev. Lett. **94**, 202002 (2005).
4. Signori, A., Bacchetta, A., Radici, M. and Schnell, G., "Investigations into the flavor dependence of partonic transverse momentum," JHEP **1311**, 194 (2013) .
5. G. Cates et al., E12-09-018, JLAB approved experiment, hallaweb.jlab.org/collab/PAC/PAC38//E12-09-018-SIDIS.pdf
6. Ciofi degli Atti, C. *et al.* "Nuclear effects in deep inelastic scattering of polarized electrons off polarized He-3 and the neutron spin structure functions," Phys. Rev. C **48**, 968 (1993).
7. Ciofi degli Atti, C., Salmè, G., and Pace, E. "Spin dependent spectral function of He-3 and the asymmetry in the process He-3 (polarized) (e (polarized),e-prime) X," Phys. Rev. C **46**, 1591 (1992) ; Kievsky, A. et al., "Neutron electromagnetic form-factors and inclusive scattering of polarized electrons by polarized He-3 and He-3 targets," Phys. Rev. C **56**, 64 (1997) .
8. Wiringa, R. B., Stoks, V. G. and Schiavilla, R. "An Accurate nucleon-nucleon potential with charge independence breaking," Phys. Rev. C **51**, 38 (1995).
9. Scopetta, S., "Neutron single spin asymmetries from semi-inclusive deep inelastic scattering off transversely polarized He-3," Phys. Rev. D **75**, 054005 (2007) .
10. Allada, K. *et al.* [JLab Hall A Collaboration], "Single Spin Asymmetries of Inclusive Hadrons Produced in Electron Scattering from a Transversely Polarized ^{3}He Target," Phys. Rev. C **89**, 042201 (2014) .
11. Kaptari, L. P., Del Dotto, A., Pace, E. Salmè, G. and Scopetta, S. "Distorted spin-dependent spectral function of an A=3 nucleus and semi-inclusive deep inelastic scattering processes," Phys. Rev. C **89**, 035206 (2014) .
12. Ciofi degli Atti, C. and Kaptari, L.P., "Semi-inclusive Deep Inelastic Scattering off Few-Nucleon Systems: Tagging the EMC Effect and Hadronization Mechanisms with Detection of Slow Recoiling Nuclei," Phys. Rev. C **83**, 044602 (2011) .
13. Del Dotto, A. *et al.*, "Flavor decomposition of transverse momentum dependent parton distributions," EPJ Web Conf. **73**, 02019 (2014).
14. Pace, E. *et al.*, "Neutron Transverse-Momentum Distributions and Polarized ^3He within Light-Front Hamiltonian Dynamics," Few Body Syst. **54**, 1079 (2013) .
15. Dirac, P.A.M., "Forms of Relativistic Dynamics", Rev. Mod. Phys. **21**, 392 (1949) .
16. Bakamjian, B. and Thomas, L. H., "Relativistic particle dynamics. 2," Phys. Rev. **92**, 1300 (1953) .
17. Scopetta, S. "Generalized parton distributions of He-3," Phys. Rev. C **70**, 015205 (2004) ; "Conventional nuclear effects on generalized parton distributions of trinucleons," Phys. Rev. C **79**, 025207 (2009) .
18. Rinaldi, M., and Scopetta, S. "Neutron orbital structure from generalized parton distributions of 3He," Phys. Rev. C **85** (2012) 062201 ; "Extracting generalized neutron parton distributions from ^3He data," Phys. Rev. C **87**, no. 3, 035208 (2013) .
19. Del Dotto, A., Pace, E., Salmè, G., and Scopetta, S., to be published.
20. C. Lorcè and B. Pasquini, "On the Origin of Model Relations among Transverse-Momentum Dependent Parton Distributions," Phys. Rev. D **84** (2011) 034039 .